

LU-ITP 2003/001  
KANAZAWA-03-03  
ITEP-LAT/2003-01  
HU-EP-02-61  
January 10, 2003

# Confinement, deconfinement and the photon propagator in 3D cQED on the lattice

M. N. Chernodub<sup>a,b,1</sup>, E.–M. Ilgenfritz<sup>c</sup> and A. Schiller<sup>d,2</sup>

<sup>a</sup> *ITEP, B. Cheremushkinskaya 25, Moscow, 117259, Russia*

<sup>b</sup> *Institute for Theoretical Physics, Kanazawa University,  
Kanazawa 920-1192, Japan*

<sup>c</sup> *Institut für Physik, Humboldt–Universität zu Berlin, D-10115 Berlin, Germany*

<sup>d</sup> *Institut für Theoretische Physik and NTZ, Universität Leipzig,  
D-04109 Leipzig, Germany*

## Abstract

We report on a lattice study of the gauge boson propagator of 3D compact QED in Landau gauge at zero and finite temperature. Non-perturbative effects are reflected by the generation of a mass  $m$  and by an anomalous dimension  $\alpha$ . These effects can be attributed to monopoles and are absent in the propagator of the regular part of the gauge field.

Three-dimensional compact electrodynamics (cQED<sub>3</sub>) shares two features with QCD, confinement and chiral symmetry breaking. Confinement of electrically charged particles is caused by a plasma of monopoles [1]. In this contribution we report how confinement is manifest in the gauge boson propagator of 3D cQED at zero and finite temperature[2, 3].

For our lattice study we have adopted the Wilson action,  $S[\theta] = \beta \sum_p (1 - \cos \theta_p)$ , where  $\theta_p$  is the field strength corresponding to the  $U(1)$  link field  $\theta_l$ . The coupling  $\beta$  is related to the lattice spacing  $a$  and the coupling  $g_3$  of the 3D continuum theory,  $\beta = 1/(a g_3^2)$ . The photon propagator is measured in Landau gauge. This gauge is defined as a global maximum of the functional  $\sum_l \cos \theta_l^G$  with respect to gauge transformations  $G$ . To select the best gauge realization, we have evaluated  $N_G$  local maxima of that functional and studied the dependence of the propagator on  $N_G$ . Simulations for  $T = 0$  have been performed on a  $32^3$  lattice, those for  $T > 0$  on a  $32^2 \times 8$  lattice. For the Monte Carlo algorithm and the gauge fixing procedure see Ref. [3]

---

<sup>1</sup>Work supported by JSPS Fellowship P01023.

<sup>2</sup>Talk presented by A.S. at Confinement 5, Gargnano, September 2002

The gauge field in lattice position space is taken as  $A_{\vec{n}+\frac{1}{2}\vec{\mu},\mu} = \sin \theta_{\vec{n},\mu}/(g_3 a)$ . Then we define the gauge boson propagator in lattice momentum space by  $D_{\mu\nu}(\vec{p}) = \langle \tilde{A}_{\vec{k},\mu} \tilde{A}_{-\vec{k},\nu} \rangle$  in terms of the Fourier transformed gauge potential for  $p_\nu = (2/a) \sin(2\pi k_\nu/L_\nu)$ . The most general tensor structure of  $D_{\mu\nu}$  at  $T = 0$  is given by

$$D_{\mu\nu}(\vec{p}) = P_{\mu\nu}(\vec{p})D(p^2) + p_\mu p_\nu F(p^2)/(p^2)^2$$

with the 3D transverse projection operator  $P_{\mu\nu}(\vec{p}) = \delta_{\mu\nu} - (p_\mu p_\nu)/p^2$ . The transverse and longitudinal propagators,  $D$  and  $F$ , are extracted by projection and are found approximately rotationally invariant or vanishing. For  $T > 0$  we have to consider two scalar functions  $D_{T/L}$  using 2D transverse/longitudinal projection operators  $P_{\mu\nu}^{T/L}$  (in 2 + 1 dimensions both are transverse):

$$D_{\mu\nu}(\vec{p}) = P_{\mu\nu}^T D_T + P_{\mu\nu}^L D_L + p_\mu p_\nu F/(p^2)^2.$$

To discuss the propagators from the monopole plasma point of view, we have decomposed the gauge fields links into singular (monopole) and regular (photon) contributions[4, 3]  $\theta_l = \theta_l^{phot} + \theta_l^{mono}$ . Once that decomposition is achieved, one is in the position to study monopole, photon and mixed contributions of  $D$  or  $D_{L/T}$  (and  $F$ ), separately.

As an example, we show in Fig. 1 (a) at  $T = 0$  the transverse and longitudinal propagators




Figure 1: (a) The propagator  $p^2 D$  and its contributions, as well as the (vanishing) propagator  $F$  vs.  $p^2$  at  $\beta = 1.8$ . (b) Fitted  $\alpha$  and  $m$  vs.  $\beta$  for  $D$ .

(multiplied by  $p^2$ ) together with the decomposition of  $D$ . The monopole part  $D^{mono}$  reaches a maximum in the low momentum region before it drops towards  $p^2 = 0$ . In contrast to the total  $D$ , the regular (photon) part is singular at  $p^2 = 0$ , and  $p^2 D^{phot}$  is flat vs.  $p^2$ .

Following Ref.[2] we describe  $D$  by the function

$$D(p^2) = (Z/\beta) m^{2\alpha}/[p^{2(1+\alpha)} + m^{2(1+\alpha)}] + C. \quad (1)$$

The photon part  $D^{phot}$  is fitted using (1) with  $\alpha = m = 0$ . At finite  $T$ , the propagator data for  $D_L$  and  $D_T$  are analyzed for  $p_3 = 0$ , as a function of  $\mathbf{p}^2$  using the same fit function (1).

The fit parameters  $\alpha$  and  $m$  of the propagators  $D$ ,  $D_L$  and  $D_T$  are presented in Figs. 1 (b) and 2 as functions of  $\beta$ . The mass  $m$  for  $T = 0$  is in good agreement with the theoretical prediction[1]




Figure 2: Fitted  $\alpha$  and  $m$  vs.  $\beta$  for  $D_L$  and  $D_T$ .

valid for a dilute monopole gas. In the confinement phase, the anomalous dimensions  $\alpha \neq 0$  for all propagators. They are functions not only of the monopole density (which is monotonously decreasing with growing  $\beta$ ). The cluster structure of the monopole configurations seems to play a significant rôle. The fit parameters  $\alpha_L$  and  $m_L$  of  $D_L$  vanish at  $\beta_c$  and beyond, a clear signal of the finite temperature phase transition[2] caused by dipole formation. Due to multiple Dirac strings wrapping around the temporal direction the behavior of  $D_T$  in the deconfined phase (at larger  $\beta$ ) is different from  $D_L$ . We carefully studied the Gribov copy dependence of the propagators. The asymptotic behaviors of  $D$  and  $D_L$  are reached quickly, contrary to the  $D_T$  propagator in the deconfinement. Further details can be found in Ref.[3]

## References

- [1] A. M. Polyakov, *Nucl. Phys.* **B120**, 429 (1977).
- [2] M. N. Chernodub, E.-M. Ilgenfritz, A. Schiller, *Phys. Rev. Lett.* **88**, 231601 (2002).
- [3] M. N. Chernodub, E.-M. Ilgenfritz, A. Schiller, *arXiv:hep-lat/0208013*, *Phys. Rev.* **D** in print.
- [4] R. J. Wensley and J. D. Stack, *Phys. Rev. Lett.* **63**, 1764 (1989).